arXiv:1302.3211v1 [math.FA] 13 Feb 2013

ON ISOMORPHISMS OF BANACH SPACES OF CONTINUOUS FUNCTIONS

GRZEGORZ PLEBANEK

Abstract. We prove that if K and L are compact spaces and $C(K)$ and $C(L)$ are isomorphic as Banach spaces then K has a π-base consisting of open sets U such that $\overline{U}$ is a continuous image of some compact subspace of L. This gives some information on isomorphic classes of the spaces of the form $C([0,1]^\kappa)$ and $C(K)$ where K is Corson compact.

1. Introduction

For a compact space K we denote by $C(K)$ the Banach space of real–valued continuous functions with the usual supremum norm. In the sequel, K and L always denote compact Hausdorff spaces.

By the classical Banach-Stone theorem, if $C(K)$ is isometric to $C(L)$ then K and L are homeomorphic, see [23], 7.8.4. On the other hand, if $C(K)$ and $C(K)$ are merely isomorphic as Banach spaces, which is denoted by $C(K) \sim C(L)$, then K and L may be far from being homeomorphic. By an important theorem due to Miljutin (see [23], 21.5.10 or [20]) for every uncountable compact metrizable space K the space $C(K)$ is isomorphic to $C[0,1]$. The Miljutin theorem, together with Bessaga and Pełczyński results from [6], gives a isomorphic classification of spaces of continuous functions on metrizable compacta.

The situation is much more complex when we try to analyse isomorphic classes of nonseparable $C(K)$ spaces. Recall for instance that there is an infinite compact space K such that $C(K)$ is indecomposable Banach space and, consequently, $C(K)$ cannot be isomorphic to $C(L)$ whenever L totally disconnected, see Koszmider [14], cf. Plebanek [21]. Such a space K has also the property that $C(K)$ is not isomorphic to $C(K+1)$ (here $K+1$ denotes the space K with one isolated point added). Another example of a space $C(K)$ such that $C(K) \sim C(L)$ for no totally disconnected compactum L has been recently found by Aviles and Koszmider [4]. It remains unclear if there are some natural examples of compact spaces K with such a property — see the problem posed by Argyros and Arvanitakis [3], page 225.

There are partial isomorphic classifications when K is a member of some concrete class of compacta, for instance dyadic spaces, see Pełczyński [20]; cf. Galego [10].

In a present paper we discuss the following problem.

2010 *Mathematics Subject Classification.* Primary 46B26, 46B03, 46E15.

Problem 1.1. *What are necessary topological relations between compact space K and L for which $C(K)$ and $C(L)$ are isomorphic as Banach spaces?*

Let $T : C(K) \to C(L)$ be an operator such that $||T|| \cdot ||T^{-1}|| < 2$. If T is surjective (so that T is an isomorphism showing that the Banach-Mazur distance between $C(K)$ and $C(L)$ is less than 2) then K is homeomorphic to L; this generalization of the Banach-Stone theorem was proved, independently, by Amir [1] and Cambern [7]. Later Jarosz [12], following Benyamini [5], showed that if such T is not necessarily onto then K is still a continuous image of a compact subspace of L. It is worth mentioning that the Amir-Cambern theorem mentioned above has been generalized beyond the context of linear mappings, see e.g. Dutreux and Kalton [9].

In [22] we have proved that if $T : C(K) \to C(L)$ is an isomorphic embedding which is moreover a positive operator then one can define a set-valued mapping from L into finite subsets of K which is surjective and upper semicontinuous. This result, partially motivated by Okuniev [19], implies that under presence of such embedding the space K inherits some topological properties of L.

We present below a certain answer to Problem 1.1 — we show that if $T : C(K) \to C(L)$ is an arbitrary isomorphism then there are many compact subspaces of K having nonempty interior that are continuous images of compact subspaces of L, see Theorem 6.1 for the precise formulation of our main result. On the way to its proof we follow some ideas from Plebanek [22]; however, dealing with arbitrary isomorphisms rather than positive embeddings seems to be much more delicate.

Theorem 6.1 in particular shows that if $C([0,1]^\kappa)$ is isomorphic to $C(L)$ then L can be continuously mapped onto $[0,1]^\kappa$ (where κ is an arbitrary cardinal number), this corollary seems to be a new result. Recall that properties of $C(K)$ for K dyadic were studied by Pełczyński [20] and Hagler [11].

When considering isomorphisms between $C(K)$ spaces we can ask which classes of compacta $\mathcal{K}$ are stable under taking such isomorphisms (that is, $K \in \mathcal{K}$ implies $L \in \mathcal{K}$ whenever $C(K) \sim C(L)$). By the classical Amir-Lindenstrauss theory, the class of Eberlein compacta (weakly compact subsets of Banach spaces) is stable in this sense, since K is Eberlein compact if and only if $C(K)$ is weakly compactly generated (see e.g. Negrepontis [18]). This motivates the following open question which has been around for several years, see e.g. 3.9 in [2], 6.45 in Negrepontis [18] or Question 1 in Koszmider [15].[1]

Problem 1.2. *Must K be Corson compact provided $C(K)$ is isomorphic to $C(L)$ where L is Corson compact?*

The class of Corson compacta is a natural extension of the class of Eberlein compact spaces. Recall that the answer to 1.2 is positive assuming Martin's axiom and the negation of continuum hypothesis, see Argyros et. al. [2]. Our main result 6.1 yields a partial positive answer to 1.2, see section 6.

---

[1] we recall some terminology in the next section

In [17] we proved that if $C(K)$ can be embedded into $C(L)$, where L is Corson compact, then K is Corson compact too provided K has some measure-theoretic property, applicable to several well-studied classes of compacta. However we were unable to verify if assuming the continuum hypothesis one can embedd $C(2^{\omega_1})$ into $C(L)$, where L is Corson compact. We now resolve this problem in the negative, see Corollary 3.5.

The paper is organized as follows. In the next section we collect some standard facts on operators and the *weak** topology in $C(K)^*$; we also recall some concepts from general topology. In section 3 and 4 we analyse the properties of the conjugate operator T^* where $T : C(K) \to C(L)$ is an arbitrary isomorphic embedding. This leads to considering finite valued maps between compact spaces, see section 4. The final section states our main result and its consequences; we also pose some related problems.

2. Preliminaries

Throughout this paper we assume that K and L denote compact Hausdorff spaces. The dual space $C(K)^*$ of the Banach space $C(K)$ is identified with $M(K)$ — the space of all signed Radon measures of finite variation; we use the symbol $M_1(K)$ to denote the unit ball of $M(K)$. Every $\mu \in M(K)$ can be written as $\mu = \mu^+ - \mu^-$ where μ^+ and μ^- are mutually singular nonnegative finite Radon measure. Recall that the variation $|\mu|$ of μ is defined as $|\mu| = \mu^+ + \mu^-$ and the natural norm in $M(K)$ is given by the formula $||\mu|| = |\mu|(K)$.

In the sequel, the space $M(K)$ is always equipped with the *weak** topology inherited from $C(K)^*$, i.e. the topology making all the functionals

$$\mu \to \int_K g \, \mathrm{d}\mu,$$

continuous, where $g \in C(K)$. Note that we usually write $\mu(g)$ for $\int_K g \, \mathrm{d}\mu$.

For any $x \in K$ we denote by $\delta_x \in P(K)$ the corresponding Dirac measure; recall that $\Delta(K) = \{\delta_x : x \in K\}$ is a subspace of $M_1(K)$ which is homeomorphic to K.

The mapping $\mu \to |\mu|$ is not *weak** continuous; nonetheless it has the following semicontinuity properties. Recall that a real-valued function φ (defined on some topological space X) is lower semicontinuous if the set $\{x \in X : \varphi(x) > r\}$ is open for every $r \in \mathbb{R}$.

Lemma 2.1. *For every compact space K*

(i) *the mapping $M(K) \ni \mu \to |\mu|(g)$ is lower semicontinuous for every nonnegative function $g \in C(K)$;*

(ii) *the mapping $M(K) \ni \mu \to |\mu|(U)$ is lower semicontinuous for every open set $U \subseteq K$.*

Proof. Let $|\mu|(g) > r$; put $\varepsilon = |\mu|(g) - r$. By regularity, there is a continuous function $h : K \to [-1, 1]$ such that $\mu(gh) > |\mu|(g) - \varepsilon/2$. Let $O(\mu)$ be an open neighbourhood of

μ defined as

$$O(\mu) = \{\nu : \nu(gh) > \mu(gh) - \varepsilon/2\}.$$

For every $\nu \in O(\mu)$ we have

$$|\nu|(g) \geq \nu(gh) > \mu(gh) - \varepsilon/2 > |\mu|(g) - \varepsilon,$$

so $\mu \in O(\mu) \subseteq \{\nu : |\nu(g)| > r\}$; this shows (i).

Part (ii) follows immediately from (i) since $|\mu|(U)$ is the supremum of $|\mu|(g)$ over all continuous $g : K \to [0,1]$ vanishing outside U. $\square$

A linear operator $T : C(K) \to C(L)$ is an isomorphic embedding if the are positive constants m_1, m_2 such that

$$m_1||g|| \leq ||Tg|| \leq m_2||g||,$$

for every $g \in C(K)$ (so that $||T|| \leq m_2$ and $||T^{-1}|| \leq 1/m_1$). When we say 'embedding' we always mean an isomorphic embedding which is not necessarily surjective.

We recall the following standard fact (see e.g. [17], Proof of 1.3).

Lemma 2.2. *If $T : C(K) \to C(L)$ is an embedding then there are a compact space L', a continuous surjection $q : L \to L'$ and an embedding $T' : C(K) \to C(L')$ such that $T'[C(K)]$ separates points of the space L'.*

With every bounded operator $T : C(K) \to C(L)$ we can associate a conjugate operator T^*, where

$$T^* : M(L) \to M(K), \quad T^*(\nu)(f) = \nu(Tf).$$

Writing $\nu_y = T^*\delta_y$ for $y \in L$, it is immediate that the mapping $L \ni y \to \nu_y \in M(K)$ is *weak** continuous. Note that $||\nu_y|| \leq ||T||$ for every $y \in L$.

A compact space K is *Corson compact* if, for some cardinal number κ, which can be taken to be equal to the topological weight of K, K is homeomorphic to a subset of the Σ–product of real lines

$$\Sigma(\mathbb{R}^\kappa) = \{x \in \mathbb{R}^\kappa : |\{\alpha : x_\alpha \neq 0\}| \leq \omega\}.$$

The class of Corson compacta has been intensively studied for its interesting topological properties and various connections to functional analysis; we refer the reader to a basic paper [2] by Argyros, Mercourakis and Negrepontis, and to extensive surveys Negrepontis [18] and Kalenda [13]. Recall that the class of Corson compacta is stable under taking closed subspaces and continuous images.

3. ISOMORPHIC EMBEDDINGS

In this section we fix a pair of compact space K and L such that there is an embedding of $C(K)$ into $C(L)$. We choose a linear operator $T : C(K) \to C(L)$ such that $||T|| = 1$ and $||T^{-1}|| = 1/m$, so that we have

$$m||g|| \leq ||Tg|| \leq ||g||,$$

for every $g \in C(K)$.

Given $y \in L$, recall that $\nu_y \in M(K)$ is the measure $\nu_y = T^*\delta_y$. In turn, for a fixed $x \in K$ there is $\mu = \mu_x \in M(L)$ such that $\mu(Tg) = g(x)$ for all $g \in C(K)$ and

$$||\mu|| = \sup\{\mu(h) : ||h|| \leq 1, h \in E\} \leq 1/m,$$

where $E = T[C(K)]$. Indeed, such a functional μ can be defined on E by the required formula and extended to $C(L)$ by the Hahn-Banach theorem.

The following fact was noted by Jarosz [12].

Lemma 3.1. *If $\mu = \mu_x$ for some fixed $x \in K$ then $||\nu_y|| \geq m$ for μ-almost all $y \in L$.*

Proof. Write again $E = T[C(K)]$ and let $N = \{y \in L : ||\delta_{y|E}|| < 1\}$.

Then $\mu(N) = 0$. Indeed, we have $N = \bigcup_{r<1} N_r$, where the sets

$$N_r = \{y \in L : ||\delta_{y|E}|| \leq r\},$$

are closed; it is therefore sufficient to check that $|\mu|(N_r) = 0$ for every $r < 1$.

Take any $\varepsilon > 0$ and $g \in C(K)$ such that $||Tg|| \leq 1$ and $\mu(Tg) > ||\mu|| - \varepsilon$. Then

$$||\mu|| - \varepsilon = |\mu|(N_r) + |\mu|(L \setminus N_r) - \varepsilon < \mu(Tg) =$$

$$= \int_{N_r} Tg \, \mathrm{d}\mu + \int_{L\setminus N_r} Tg \, \mathrm{d}\mu \leq r|\mu|(L) + |\mu|(L \setminus N_r),$$

which gives $|\mu|(N_r) \leq \varepsilon/(1 - r)$ and, consequently, $|\mu|(N_r) = 0$, as $\varepsilon > 0$ is arbitrary.

Now for any $y \in L \setminus N$ and any $\varepsilon' > 0$ there is $g \in C(K)$, $||Tg|| \leq 1$ such that $|Tg(y)| > 1 - \varepsilon'$. Then $||g|| \leq 1/m$ and $|\nu_y(g)| = |Tg(y)| > 1 - \varepsilon'$. This implies $||\nu_y|| \geq m$, and we are done. □

Lemma 3.2. *Consider a fixed $x \in K$ and the measure $\mu = \mu_x$. Take any $m' < m$ and ε such that $0 < \varepsilon < (m - m')/2$, and suppose that $\widehat{L}$ is a compact subset of L with the following properties:*

(i) *$||\nu_y|| \geq m$ for every $y \in \widehat{L}$;*
(ii) *the function $\widehat{L} \ni y \to ||\nu_y||$ has oscillation $\leq \varepsilon$ at each point of $\widehat{L}$;*
(iii) *$|\mu|(L \setminus \widehat{L}) < \varepsilon$.*

Then there is $y \in \widehat{L}$ such that $|\nu_y\{x\}| \geq m'$.

Proof. Let us write $m'' = (m+m')/2$.

Let H be any neighbourhood of x and $f_H : K \to [0,1]$ be a continuous function such that $f_H(x) = 1$ and $f_H = 0$ outside H.

CLAIM. There is $y \in \widehat{L}$ such that $Tf_H(y) > m''$.

Indeed, otherwise

$$1 = f_H(x) = \mu(Tf_H) = \int_{\widehat{L}} Tf_H \, \mathrm{d}\mu + \int_{L\setminus\widehat{L}} Tf_H \, \mathrm{d}\mu \leq$$

$$\leq m''|\mu|(\widehat{L}) + \varepsilon \leq \frac{m''}{m} + \varepsilon,$$

since $|\mu|(\widehat{L}) \leq |\mu|(L) = ||\mu|| \leq 1/m$. It follows that

$$\varepsilon \geq 1 - m''/m = \frac{m-m''}{m} = \frac{m-m'}{2m} \geq \frac{m-m'}{2},$$

a contradiction with our choice of ε; this verifies Claim.

Let $\mathcal{H}(x)$ be a local base at x; for every $H \in \mathcal{H}(x)$ we choose a continuous function $f_H : K \to [0,1]$, such that $f_H(x) = 1$, $f_H = 0$ on H^c. By Claim there is $y_H \in L_i$ such that $\nu_{y_H}(f_H) > m''$.

The points y_H form a net indexed by $H \in \mathcal{H}(x)$ ordered by inverse inclusion. Passing to a finer subnet if necessary we can assume that y_H converge to some $y \in L$; then $y \in \widehat{L}$ since $\widehat{L}$ is compact. We shall check that ν_y is as required. By regularity, in order to verify $|\nu_y|(\{x\}) \geq m'$ it suffices to check that $|\nu_y|(U) \geq m'$ for every open set $U \ni x$.

Given an open set $U \ni x$, choose a continuous function $g : K \to [0,1]$ such that $g = 0$ outside U and $g = 1$ on an open set V containing x. For any $H \in \mathcal{H}(x)$ with $H \subseteq V$ we have $|\nu_{y_H}|(g) \geq |\nu_{y_H}|(f_H) > m''$. Moreover, it follows from (ii) that $||\nu_{y_H}|| < ||\nu_y|| + \varepsilon$ for cofinally many H. Hence

$$|\nu_{y_H}|(1-g) < |\nu_{y_H}|(K) - m'' \leq |\nu_y|(K) + \varepsilon - m'' < |\nu_y|(K) - m',$$

because $\varepsilon - m'' < -m'$ by our assumption on ε. Since $\nu_H \to \nu_y$, it follows from Lemma 2.1(i) and the above inequality that

$$|\nu_y|(1-g) \leq |\nu_y|(K) - m'.$$

We conclude that $|\nu_y|(U) \geq |\nu_y|(g) \geq m'$ and the proof is complete. □

Proposition 3.3. *If $T : C(K) \to C(L)$ is an isomorphic embedding then, writing $\nu_y = T^*\delta_y$ for $y \in L$, for every $x \in K$ we have*

$$\sup\{|\nu_y(\{x\})| : y \in L\} \geq \frac{1}{||T||||T^{-1}||}.$$

Proof. Clearly we can assume that $||T|| = 1$ and denote $m = 1/||T^{-1}|| > 0$. The function $L \ni y \to ||\nu_y||$ is lower semicontinuous hence Borel. Given $x \in K$, by Lemma 3.1 $||\nu_y|| \geq m$ μ_x-almost everywhere. By the Lusin theorem we can therefore find for any $m' < m$ and any $\varepsilon > 0$ a compact set $\widehat{L} \subseteq L$ with $|\mu|(L \setminus \widehat{L}) < \varepsilon$ and such that $\widehat{L} \ni y \to ||\nu_y||$ is continuous. Applying Lemma 3.2 we conclude the proof. $\square$

Recall that a set $L \subseteq M(K)$ is said to be *m-norming*, where $m > 0$, if for every $g \in C(K)$ there is $\mu \in L$ such that $|\mu(g)| \geq m \cdot ||g||$.

Corollary 3.4. *If K is a compact space and $L \subseteq M_1(K)$ is weak* compact m-norming set then* $\sup\{|\nu|(\{x\}) : \nu \in L\} \geq m$ *for every $x \in K$.*

Proof. Since L is *weak** compact it is bounded in $M(K)$. For such a space L the mapping $T : C(K) \to C(L)$ given by $Tg(\nu) = g(\nu)$ is a bounded operator and T is an embedding as $||T^{-1}|| \leq 1/m$. Hence the assertion follows from Proposition 3.3. $\square$

We conclude this section by showing the following result on Corson compacta.

Corollary 3.5. *Let K be such a compact space that* $\mathrm{card}K > \mathfrak{c}$ *and* $\mathrm{card}[C(K)] = \mathfrak{c}$. *Then $C(K)$ cannot be isomorphically embedded into $C(L)$ with L being Corson compact.*

Proof. Suppose otherwise and let $T : C(K) \to C(L)$ be an embedding, where L is Corson compact. Since the class of Corson compacta is closed under taking continuous images, it follows from Lemma 2.2 that we can additionally assume that the functions from $T[C(K)]$ distinguish points of L. As the space $C(K)$ has cardinality $\mathfrak{c}$, it follows that there are $\mathfrak{c}$ many continuous functions on L separating points; this implies that the topological weight of L does not exceed $\mathfrak{c}$. Then L is homeomorphic to a subspace of $\Sigma(\mathbb{R}^{\mathfrak{c}})$ (which is of cardinality $\mathfrak{c}$) so in particular $\mathrm{card}L \leq \mathfrak{c}$.

On the other hand, taking $0 < r < 1/(||T|| \cdot ||T^{-1}||)$, by Proposition 3.3 to every $x \in K$ we can assign $\theta(x) \in L$ such that $|\nu_{\theta(x)}(\{x\})| \geq r$. Since every $|\nu_y|$ is a finite measure there may be only finite number of atoms of measure $|\nu_y|$ greater than r. It follows that $\theta : K \to L$ is a finite-to-one map and therefore $\mathrm{card}K \leq \mathfrak{c}$, contrary to our assumption. $\square$

At this point it is worth recalling that assuming Martin's axion and the negation of the continuum hypothesis (MA+ ¬CH for short), if L is Corson compact then $M_1(L)$ is also Corson compact in its *weak** topology, Argyros et al. [2]. Consequently, if $T : C(K) \to C(L)$ is an embedding then $T^*[M_1(L)]$ is Corson compact and so is K ($T^*[M_1(L)]$ contains a ball in $M(K)$ because T^* is onto and K can be embedded into the space of measures via the mapping $K \ni x \to \delta_x$).

The above argument fails if CH is assumed but using Corollary 3.5 we can prove the following.

Corollary 3.6. *Assuming CH, $C(2^{\omega_1})$ cannot be embedded into $C(L)$ with L being Corson compact.*

Proof. Under CH the Cantor cube $K = 2^{\omega_1}$ has cardinality $2^{\mathfrak{c}} > \mathfrak{c}$. Moreover, there are only $\mathfrak{c}$ many continuous functions on K because every such a function is determined by countably many coordinates. Hence we can apply Corollary 3.5. $\square$

We do not know if Corollary 3.6 can be proved without any extra set-theoretic axioms, i.e. we do not know what happens when $2^{\omega_1} = \mathfrak{c}$ but MA does not hold. It will become clear in section 6 that spaces such as $C(2^{\omega_1})$ cannot be isomorphic to a space $C(L)$ whenever L is Corson compact.

4. ISOMORPHISMS

We now consider a fixed pair of compact spaces K and L and an isomorpism T from $C(K)$ **onto** $C(L)$. As before, we usually assume that $||T|| = 1$ and $||T^{-1}|| = 1/m$. Now for every $x \in K$ the measure μ_x on L is uniquely defined by the formula $\mu_x(Tg) = g(x)$. Moreover, $\mu_x = (T^{-1})^*(\delta_x)$ so the mapping $x \to \mu_x$ is *weak** continuous. Note that $1 \leq ||\mu_x|| \leq 1/m$ for every $x \in K$ and $m \leq ||\nu_y|| \leq 1$ for every $y \in L$.

In the sequel, we say that a set $G \subseteq K$ is half-open if $G = C \cap V$ for some closed $C \subseteq K$ and some open $V \subseteq K$.

Lemma 4.1. *For every $\varepsilon > 0$ and any closed $Z \subseteq L$ there is a finite partition $\{G_j : j \leq j_0\}$ of K into half-open sets G_j such that for every $j \leq j_0$ and $x_1, x_2 \in G_j$ we have*

$$(*) \quad \big||\mu_{x_1}|(Z) - |\mu_{x_2}|(Z)\big| < \varepsilon.$$

Proof. As the function $x \to ||\mu_x||$ is lower semicontinuous, the sets

$$K_i = \{x \in K : ||\mu_x|| \leq 1 + i(\varepsilon/2)\},$$

are closed and $K_i = K$ for i big enough. To prove the lemma it will do to show that every set of the form $G = K_{i+1} \setminus K_i$ can be decomposed into finitely many half-open sets G_j so that (*) holds for any $x_1, x_2 \in G_j$.

Again, since $x \to |\mu_x|(L \setminus Z)$ is lower semicontinuous by Lemma 2.1(ii), the sets

$$F_j = \{x \in K : |\mu_x|(L \setminus Z) \leq j(\varepsilon/2\},$$

are closed. Now if we put $G_j = G \cap (F_{j+1} \setminus F_j)$ then for $x_1, x_2 \in G_j$ we have

$$\big||\mu_{x_1}|(Z) - |\mu_{x_2}|(Z)\big| \leq \big||\mu_{x_1}|(L) - |\mu_{x_2}|(L)\big| + \big||\mu_{x_1}|(L \setminus Z) - |\mu_{x_2}|(L \setminus Z)\big| < \varepsilon.$$

$\square$

Lemma 4.2. *If $\{G_j : j \leq j_0\}$ is a partition of K into half-open sets then for every nonempty open set $U \subseteq K$ there is nonempty $V \subseteq U$ such that $V \subseteq G_j$ for some $j \leq j_0$.*

Proof. Let us write $G_j = V_j \cap C_j$, where V_i is open and C_j is closed. Note that there is $j \leq j_0$ such that $U \cap V_j \cap \mathrm{int}(C_j) \neq \emptyset$ for, otherwise, we would have

$$U \subseteq \bigcup_{j \leq j_0} (C_j \setminus \mathrm{int}(C_j),$$

but U cannot be covered by a finite number of nowhere dense sets. $\square$

Lemma 4.3. *Let $Z_1 \subseteq Z_2 \subseteq Z_3 = L$ be closed subsets of L and let $\eta > 0$. Suppose that $U \subseteq K$ is a neighbourhood of $x_0 \in K$ such that for $l = 1, 2, 3$ and every $x \in U$*

$$(*) \quad \big||\mu_x|(Z_l) - |\mu_{x_0}|(Z_l)\big| < \eta.$$

Then there is an open set V with $x_0 \in V \subseteq U$ and a closed set $L_1 \subseteq Z_2 \setminus Z_1$ such that for every $x \in V$ we have

$$|\mu_x|(L_1) > |\mu_x|(Z_2 \setminus Z_1) - 4\eta.$$

Proof. Note that there is a continuous function $h : L \to [0, 1]$ such that its support $S = \overline{\{y \in L : h(y) > 0\}}$ is disjoint from Z_1 and

$$\mu_{x_0}(h) > |\mu_{x_0}|(L \setminus Z_1) - \eta.$$

We put $L_1 = S \cap Z_2$ and define V as

$$V = \{x \in U : \big|\mu_x(h) - \mu_{x_0}(h)\big| < \eta\};$$

then V is open since the mapping $x \to \mu_x(h)$ is continuous.

Now for any $x \in V$

$$|\mu_x|(S) \geq \mu_x(h) > \mu_{x_0}(h) - \eta > |\mu_{x_0}|(L \setminus Z_1) - 2\eta > |\mu_x|(L \setminus Z_1) - 4\eta,$$

where the last inequality follows from (*) applied to $l = 1$ and $l = 3$. It follows that

$$|\mu_x|(L_1) \geq |\mu_x|(Z_2 \setminus Z_1) - 4\eta,$$

for $x \in V$, and we are done. $\square$

Theorem 4.4. *Suppose that K and L are compact spaces such that $C(K)$ is isomorphic to $C(L)$; let $T : C(K) \to C(L)$ be an isomorphism such that $||T|| = 1$, $||T^{-1}|| = 1/m$. Then, given $m' < m$ and $\varepsilon > 0$, for every nonempty open set $U \subseteq K$ there is a nonempty open set $V \subseteq U$ and a compact set $\widehat{L} \subseteq L$ such that*

(a) *the function $\widehat{L} \ni y \to ||\nu_y||$ has oscillation smaller than ε at each $y \in \widehat{L}$;*
(b) *for every $x \in V$ there is $y \in \widehat{L}$ such that $|\nu_y(\{x\})| \geq m'$.*

Proof. Let $Z_i = \{y \in L : ||\nu_y|| \leq m + \varepsilon i\}$. Then $Z_1 \subseteq Z_2 \subseteq \ldots \subseteq L$ are closed and $Z_{i_0} = L$ for some i_0. Take $\eta > 0$ such that $\eta < \varepsilon/(4i_0)$.

Applying Lemma 4.1 to each Z_i we infer that K can be partitioned into a finite number of half-open sets G_j, $j \leq j_0$, such that for every j and $x, x' \in G_j$ we have

$$\big||\mu_x|(Z_i) - |\mu_{x'}|(Z_i)\big| < \eta,$$

for all $i \leq i_0$.

Given open $\emptyset \neq U \subseteq K$, by Lemma 4.2 there is $\emptyset \neq W \subseteq U$ such that $W \subseteq G_j$ for some $j \leq j_0$. Take $x_0 \in W$; by Lemma 4.3 (applied a finite number of times) there are an open set V, where $x_0 \in V \subseteq W$, and closed sets $L_i \subseteq Z_{i+1} \setminus Z_i$ such that for every $x \in V$ and every $i \le i_0$

$$|\mu_x|(L_i) \ge |\mu_x|(Z_{i+1} \setminus Z_i) - 4\eta.$$

Letting $\widehat{L} = \bigcup_{i \le i_0} L_i$ we have $|\mu_x|(\widehat{L}) > |\mu_x|(L) - \varepsilon$ for every $x \in V$.

We finish the proof referring to Lemma 3.2. Indeed, we can of course assume that $\varepsilon < (m - m')/2$; the set $\widehat{L}$ satisfies (ii) of Lemma 3.2 since L_i are pairwise disjoint. $\square$

5. FINITE VALUED MAPS

Let K and L be compact spaces. We shall consider in this section finite-valued set mappings form L into $[K]^{<\omega}$, the family of all finite subsets of K.

Definition 5.1. A function $\varphi : L \to [K]^{<\omega}$ is said to be

(i) upper semicontinuous if the set $\{y \in L : \varphi(y) \subseteq U\}$ is open for every open $U \subseteq K$;

(ii) lower semicontinuous if the set $\{y \in L : \varphi(y) \cap U \neq \emptyset\}$ is open for every open $U \subseteq K$;

(iii) surjective if $\bigcup_{y \in L} \varphi(y) = K$.

It is clear that both upper and lower semicontinuity imply continuity in case the set valued function is single-valued. The function that we consider below satisfy a certain ε-version of upper semicontinuity and are, in a sense, locally lower semicontinuous.

Let us note that if K is the space $[0,1] \times \{0,1\}$ with the topology defined by the lexicographic order (that is K is the familiar double arrow space with two additional isolated points) then there is a surjection $\varphi : [0,1] \to [K]^2$ which is upper semicontinuous while no essential part of K is a continuous image of the subset of $[0,1]$.

We consider set valued mappings that arise in connection with isomorphisms $T : C(K) \to C(L)$, as described in Theorem 4.4. Let us briefly summarize the context.

5.2 Throughout this section we assume that we are given two compact spaces K and L, and a mapping $L \ni y \to \nu_y \in M_1(K)$ such that for some $m > 0$ (previously called m')

5.2(a) $y \to \nu_y$ is *weak** continuous;
5.2(b) $y \to |\nu_y|(U)$ is lower semicontinuous for every open $U \subseteq K$;
5.2(c) for every $x \in K$ there is $y \in L$ with $|\nu_y(\{x\})| \ge m$.
5.2(d) the mapping $y \to ||\nu_y||$ has oscillation $< \varepsilon$ at every $y \in L$.

Here ε is a positive constant depending only on m; it will turn out that any $\varepsilon < m/18$ will do.

For any $r > 0$ we define

$$\varphi_r(y) = \{x \in K : |\nu_y(\{x\})| \ge r\}.$$

Then $\varphi_r : L \to [K]^{\leq p}$ where p is the integer part of $1/r$ since $||\nu_y|| \leq 1$ for every $y \in L$. Note also that for every $r \leq m$, φ_r is onto by 5.2(c).

Lemma 5.3. *If $U \subseteq K$ is open and $\varphi_{r-\varepsilon}(y) \subseteq U$ for some $y \in L$ then there is a neighbourhood W of y such that $\varphi_r(z) \subseteq U$ for every $z \in W$.*

Proof. As $\varphi_{r-\varepsilon}(y) \subseteq U$, for any $x \in K \setminus U$ we have $\left|\nu_y(\{x\})\right| < r - \varepsilon$ and therefore there is an open set $U_x \ni x$ such that $|\nu_y|(\overline{U_x}) < r - \varepsilon$. There are x_i, $i \leq i_0$, such that the sets $U_i = U_{x_i}$ form a finite cover of $K \setminus U$. Let

$$\eta = \min\{r - \varepsilon - |\nu_y|(\overline{U_i}) : i \leq i_0\}.$$

Using 5.2(b) we can find an open set $W \ni y$ such that if $z \in W$ then

$$|\nu_z|(K \setminus \overline{U_i}) > |\nu_y|K \setminus \overline{U_i}) - \eta,$$

for every $i \leq i_0$. By 5.2(d) we can moreover assume that $||\nu_z|| < ||\nu_y|| + \varepsilon$ for $z \in W$.

Take any $z \in W$ and $x \in K \setminus U$. Then $x \in U_i$ for some $i \leq i_0$ and hence

$$|\nu_z|(\{x\}) \leq |\nu_z|(\overline{U_i}) = |\nu_z|(K) - |\nu_z|(K \setminus \overline{U_i}) <$$

$$< |\nu_y|(K) + \varepsilon - |\nu_y|(K \setminus \overline{U_i}) + \eta = |\nu_y|(\overline{U_i}) + \eta + \varepsilon \leq r.$$

Hence $\varphi_r(z) \subseteq U$ for $z \in W$, and we are done. □

Lemma 5.4. *For any $r > \varepsilon$*

$$\overline{\bigcup_{y \in L} \varphi_r(y)} \subseteq \bigcup_{y \in L} \varphi_{r-\varepsilon}(y).$$

Proof. Suppose that $x \notin \varphi_{r-\varepsilon}(y)$ for every $y \in L$. Then for every $y \in L$ there is $U_y \ni x$ such that $\varphi_{r-\varepsilon}(y) \subseteq K \setminus \overline{U_y}$. By Lemma 5.3 $\varphi_r(z) \subseteq K \setminus \overline{U_y}$ for z from some open set $V_y \ni y$. Take a finite cover V_{y_j}, $j \leq j_0$ of L and let $U = \bigcap_{j \leq j_0} U_{y_j}$. Then U is disjoint from $\bigcup_{y \in L} \varphi_r(y)$, so $x \notin \overline{\bigcup_{y \in L} \varphi_r(y)}$, as required. □

Proposition 5.5. *Let K and L be compact spaces such that there is a mapping $L \ni y \to \nu_y \in M_1(K)$ as in 5.2. Then inside every nonempty open subset of K there exists a compact set $\widehat{K}$ with nonempty interior which is a continuous image of some compact set $L_1 \subseteq L$.*

Proof. We shall find a set $\widehat{K} \subseteq K$ with the required property; it will become clear how to modify the argument to get a set $\widehat{K}$ that is contained in a given open nonempty subset of K.

We shall first reduce the prove to the case in which for some r

$$(\dagger) \quad \bigcup_{y \in L} \varphi_r(y) = K; \text{ while}$$

$$(\ddagger) \varphi_{(3/2)r}(y) = \emptyset \text{ for every } y \in L.$$

Such a reduction may be done as follows: Let

$$r_0 = \inf\{r > 0 : \operatorname{int}\left(\overline{\bigcup_{y\in L} \varphi_r(y)}\right) = \emptyset\}.$$

Then $m \leq r_0 \leq 1$. Choose r_1 such that $r_1 + \varepsilon < r_0 < (3/2)r_1$. By our choice of constants there is a compact set K_1 with nonempty interior such that

$$K_1 \subseteq \overline{\bigcup_{y\in L} \varphi_{r_1+\varepsilon}(y)} \setminus \overline{\bigcup_{y\in L} \varphi_{(3/2)r_1}(y)}.$$

Then $K_1 \subseteq \bigcup_{y\in L} \varphi_{r_1}(y)$ by Lemma 5.4 and $|\nu_y|(\{x\}) < (3/2)r_1$ for every $x \in K_1$ and $y \in L$.

Now we might consider K_1 instead of K from this point; instead let us simply assume that (†) and ‡) are satisfied (and r is fixed).

For the sake of the proof of Claim below we need to know that $(3/2)r + 2\varepsilon < 2(r - 2\varepsilon)$. This is equivalent to $\varepsilon < r/12$. Since $r \geq (2/3)m$ we can fix ε to be any positive constant $< m/18$.

CLAIM. Any $x \in K$ has an open neighbourhood U such that $\operatorname{card}(\varphi_{r-2\varepsilon}(y) \cap U) \leq 1$ for every $y \in L$.

To verify Claim take any $y \in L$ and choose $U_y \ni x$ such that $|\nu_y|(\overline{U_y}) < (3/2)r$. Then by Lemma 2.1(ii) there is open $V_y \ni y$ such that for every $z \in V$

$$|\nu_y|(K) > |\nu_z|(K) - \varepsilon;$$

$$|\nu_z|(K \setminus \overline{U_y}) > |\nu_y|(K \setminus \overline{U_y}) - \varepsilon.$$

The first requirement can be fulfilled by 5.2(d) while the second by Lemma 2.1(ii). It follows that for any $z \in V_y$ we have

$$|\nu_z|(\overline{U_y}) = |\nu_z|(K) - |\nu_z|(K \setminus \overline{U_y}) \leq$$

$$\leq |\nu_y|(K) - |\nu_y|(K \setminus \overline{U_y}) + 2\varepsilon = |\nu_y|(\overline{U_y}) + 2\varepsilon < (3/2)r + 2\varepsilon.$$

As we have fixed ε so that $(3/2)r + 2\varepsilon < 2(r - 2\varepsilon)$ (see the remark prior to Claim) then it follows that $\varphi_{r-\varepsilon}(y)$ cannot intersect $\overline{U}$ at two points.

Take a finite set $s \subseteq L$ such that the sets V_y, for $y \in s$ form a cover of L. Then $U = \bigcap_{y\in s} U_y$ is as required; this finishes the proof of Claim.

We now let

$$Y = \{y \in L : \varphi_r(y) \cap \overline{U} \neq \emptyset\},$$

where U is the set as in Claim. Then $\varphi_{r-\varepsilon}(y) \cap \overline{U} \neq \emptyset$ for every $y \in \overline{Y}$ by Lemma 5.3.

Finally, we define $h : \overline{Y} \to \overline{U}$ so that $h(y)$ is the unique point in $\overline{U} \cap \varphi_{r-\varepsilon}(y)$. Then h maps $\overline{Y}$ onto $\overline{U}$ so it remains to check that h is continuous.

Let the set $F \subseteq \overline{U}$ be closed. Then $h^{-1}[F] = A$, where

$$A = \{y \in \overline{Y} : \varphi_{r-\varepsilon}(y) \cap F \neq \emptyset\},$$

and A is closed. Indeed, if $y \notin A$, i.e. $\varphi_{r-\varepsilon} \cap F = \emptyset$ then $\varphi_{r-2\varepsilon}(y) \cap F = \emptyset$ as well, and it follows from Lemma 5.3 that $\varphi_{r-\varepsilon}(z) = \emptyset$ for all z from some neighbourhood of z. ☐

6. RESULTS AND QUESTIONS

Now we are ready to state and prove our main result.

Theorem 6.1. *Let K and L be compact spaces such that $C(K)$ is isomorphic to $C(L)$. Then for every nonempty open set $U \subseteq K$ there exists a nonempty open set $V \subseteq U$ such that $\overline{V}$ is a continuous image of some compact subspace of L.*

Proof. Apply Theorem 4.4 for any positive $m' < m$; this gives $V \subseteq K$ and $\widehat{L} \subseteq L$ satisfying (a) and (b) of Theorem 4.4. Then use Proposition 5.5 with $\overline{V}$ playing the role of K, $\widehat{L}$ replacing L, and m' standing for m (with any ε such that $0 < \varepsilon < m'/12$). ☐

Of course the above theorem gives no information for spaces K having dense sets of isolated points. On the other hand, the result has the following consequences.

Corollary 6.2. *Suppose that K has a π-base consisting of open sets $V \subseteq K$ such that K is a continuous image of $\overline{V}$. Then whenever L is a compact space such that $C(K) \sim C(L)$ then K is a continuous image of a compact subspace of L.*

Corollary 6.3. *Given a cardinal number κ and a compact space L, if $C[0,1]^\kappa$ is isomorphic to $C(L)$ then L maps continuously onto $[0,1]^\kappa$.*

Proof. Clearly $[0,1]^\kappa$ satisfies the assumption of Corollary 6.2 so $C(K) \sim C(L)$ implies that there is a compact subspace $L_1 \subseteq L$ and a continuous surjection $\theta : L_1 \to [0,1]^\kappa$; in turn such θ can be extended to a continuous mapping $L \to [0,1]^\kappa$ by the Tietze extension theorem. ☐

Corollary 6.4. *If L is Corson compact and $C(K) \sim C(L)$ for some compact K then K has a π-base of sets with Corson compact closures. In particular, K is Corson compact itself whenever K is homogeneous.*

Proof. The first assertion follows from Theorem 6.1 since the class of Corson compacta is closed under taking compact subspaces and continuous images.

If K is homogeneous that it follows that can be covered by a finite family $\{V_i : i \leq i_0\}$ of open sets where $\overline{V_i}$ is Corson compact for every i. Then a disjoint union $K' = \bigoplus_{i \leq i_0} \overline{V_i}$ is Corson compact and so is K since K is a continuous image of K'. ☐

In connection with Problem 1.2 which, in full generality remains open, one can ask the following questions.

Problem 6.5. *Does $C(K) \sim C(L)$ imply that K and L have the same tightness?*

Must K be a Frechet space (sequentially compact) whenever $C(K) \sim C(L)$ with L being Frechet (sequentially compact)?

Recall that the tightness of K is $\leq \kappa$ if for every $A \subseteq K$ and $x \in \overline{A}$ there is A_0 of cardinality $\leq \kappa$ such that $x \in \overline{A_0}$. K is a Frechet space if $x \in \overline{A}$ implies that some sequence from A converges to x. Every Corson compactum has countable tightness and is a Frechet space.

The Frechet property is a generalization of first-countability. It is worth noting that there is a first-countable compact space L and a compact space K such that $C(K) \sim C(L)$ but K is not first-countable; this was comunicated to us by Koszmider [16].

We have proved in [22] that if there is a positive embedding of $C(K)$ into $C(L)$ then the tightness of K is less than or equal to the tightness of L and, K is Frechet (sequentially compact) whenever L is.

Finally, let us remark that the main obstacle in analysing embeddings (or isomorphisms) $T : C(K) \to C(L)$ is that the mapping $L \ni y \to ||T^*\delta_y||$ need not to be continuous. Assuming that such a mapping is actually continuous one can show that there is a positive embedding $C(K) \to C(L \times 2)$, see [22].

Acknowledgement. The author was partially suported by by MNiSW Grant N N201 418939 (2010–2013).

References

[1] D. Amir, *On isomorphis of continuous function spaces*, Israel J. Math, 3 (1965), 205-210.

[2] S. Argyros, S. Mercourakis, S. Negrepontis, *Functional analytic properties of Corson–compact spaces*, Studia Math. 89 (1988) 197–228.

[3] S. Argyros, A.D. Arvanitakis, *A characterization of regular averaging operators and its consequences*, Studia Math. 151 (2002), 207–226.

[4] A. Aviles, P. Koszmider, *S continuous image of a Radon-Nikodym compact space which is not Raon-Nikodym*, preprint (2011).

[5] Y. Benyamini, *Small into–isomorphisms between spaces of continuous functions*, Proc. Amer. Math. Soc. 83 (1981), 479–485.

[6] Cz. Bessaga, A. Pełczyński, Spaces of continuous functions IV (On isomorhic classification of $C(S)$ spaces), Studia Math. 19 (1960), 53–62.

[7] M. Cambern, *On isomorphisms with small bounds*, Proc. Amer. Math. Soc. 18 (1967), 1062–1066.

[8] J. Diestel, *Sequences and series in Banach spaces*, Springer 1984.

[9] Y. Dutrieux, N.J. Kalton, *Perturbations of isometries between $C(K)$-spaces* Studia Math. 166 (2005), 181–197.

[10] E. Medina Galego, *On isomorphic classification of* $C(\mathbf{2}^{\mathrm{m}} \oplus [0,\alpha])$, Fund. Math. 204 (2009), 87–95.

[11] J. Hagler, *On the structure of S and $C(S)$ for S dyadic*, Trans. Amer. Math. Soc. 214 (1975), 415–428

[12] K. Jarosz, *Into isomorphisms of spaces of continuous functions*, Proc. Amer. Math. Soc. 90 (1984), 373–377.

[13] O. Kalenda, *Valdivia compact spaces in topology and Banach space theory*, Extracta Math. 15 (2000), 1–85.

[14] P. Koszmider, *A Banach space of continuous functions with few operators*, Math. Ann. 330 (2004), 151–183.

[15] P. Koszmider, *The interplay between compact spaces and the Banch spaces of their continuous functions*, in: Open Problems in Topology II, E. Pearl (edt.), Elsevier (2007).

[16] P. Koszmider, Note of March, 2009.

[17] W. Marciszewski, G. Plebanek, *On Corson compacta and embeddings of C(K) spaces*, Proc. Amer. Math. Soc. 138 (2010), 4281-4289.

[18] S. Negrepontis, *Banach spaces and topology*, in: Open problems in topology, J. van Mill, G.M. Reed (edts.), Chapter 23.

[19] O. Okunev, *Tightness of compact spaces is preserved by the t-equivalence relation*, Com. Math. Univ. Car. 43 (2002), 335-342.

[20] A. Pełczyński, *Linear extensions, linear averagins and their applications to linear topological characterizations of spaces of continuous functions*, Dissertationes Math. 58 (1968).

[21] G. Plebanek, *A construction of a Banach space $C(K)$ with few operators*, Topology Appl. 143 (2004), 217–239.

[22] G. Plebanek, *On positive embeddings of $C(K)$ spaces*, preprint (2012).

[23] Z. Semadeni, *Banach spaces of continuous functions*, PWN (1971).

INSTYTUT MATEMATYCZNY, UNIWERSYTET WROCŁAWSKI
E-mail address: `grzes@math.uni.wroc.pl`